

Evaluation of the Compulsory Drug Treatment Program: Within-treatment change

Sasha Nahleen & Mark Howard

Aim

To examine whether participation in the Compulsory Drug Treatment Program (CDTP) was associated with change in risk factors that are relevant to the drug-crime cycle.

Method

CDTP participants completed assessments in a battery of psychometric measures before starting the program and again after each of the three program stages. The sample included participants who completed at least one of these assessments ($n = 538$). Within-treatment change on measures was calculated using mixed effects models with program stage as a fixed factor and participants as a random factor.

Results

On average, participants showed significant within-treatment improvement on key treatment targets including self-efficacy, quality of life, attitudes towards crime and offending, thinking styles that support criminality, problem solving ability, psychological and social functioning, and self-control. Results also showed incremental improvements after each of the program stages. Most measures on key responsivity factors showed that treatment readiness was high, and perceived coercion low, before starting the program and remained stable over time.

Conclusion

This study is one of the first to evaluate intermediate processes of change associated with participation in the CDTP, and gives promising initial indications of improvements in a range of treatment targets that are relevant to drug-related offending. Additional planned evaluations will be beneficial to further assess the processes and outcomes associated with this important judicial avenue for addressing substance use-related needs among people involved in the criminal justice system.

INTRODUCTION

A history of drug use is common among people entering prison. Survey results found that 65% of prison entrants across Australia reported using illicit drugs during the previous 12 months (Australian Institute of Health and Welfare, 2019) and a urinalysis study found that the majority of police detainees (82%) tested positive to at least one drug type (Voce & Sullivan, 2020). One meta-analysis of 30 studies showed that the odds of offending were three to four times greater for drug users than non-drug users (Bennett et al., 2008) and another confirmed the overall predictive relationship between substance abuse and reoffending (Dowden & Brown, 2002).

Given the extent and risk-relevance of drug use among prison populations, various alcohol and drug interventions are well embedded into criminal justice pathways. One such example is the Compulsory Drug Treatment Program (CDTP) in NSW, which is designed to divert recidivist, drug-dependent offenders away from the traditional criminal justice system and break the drug-crime cycle with compulsory treatment and rehabilitation. There are four intended legislative objectives of the program: 1) to provide a comprehensive program of compulsory treatment and rehabilitation for drug dependent people who repeatedly resort to criminal activity to support their dependency; 2) to effectively treat drug dependency, eliminating illicit drug use while in the program and reducing the likelihood of relapse on release, 3) to promote the re-integration of participants into the community, and 4) to prevent and reduce crime by reducing the need to resort to criminal activity to support dependency (Crimes (Administration of Sentences) Act 1999, s. 106B).

During the program, offender rehabilitation integrates both the risk-need-responsivity model (RNR; Andrews & Bonta, 2010) and the good lives

model (Ward, 2002; see Birgden & Grant, 2010 for more detail). The RNR model focuses on addressing and managing dynamic risk factors for drug-related offending (e.g., substance use and impulsivity) through targeted treatment. In contrast, the good lives model supports the development of a "good life" and guides treatment by emphasizing fulfilling individuals' basic human needs through pro-social means, promoting autonomy, and a therapeutic alliance between staff and offenders. Together, these models ensure that participants receive comprehensive support for both risk management and improved well-being, ultimately facilitating community reintegration.

Only sentenced individuals who might be eligible for the program are referred to the Drug Court for an eligibility and suitability assessment. If the defendant is considered eligible and suitable, the Drug Court imposes a Compulsory Drug Treatment Order. The alternative to this Order is serving the full term of the sentence in a mainstream prison. Neither the Crown nor the offender has a right to object to, or appeal against, the referral to the Drug Court for consideration of an Order. Program participants are housed in the purpose-built Compulsory Drug Treatment Correctional Centre (CDTCC) which opened in 2006. Participants enter into a Personal Plan agreement which identifies their dynamic risk factors, outlines treatment conditions, and specifies the rewards for meeting these conditions and the sanctions for not meeting them. They then move through three stages of treatment, each of which last for at least six months. Successfully meeting the conditions of the Personal Plan is rewarded with progression toward community reintegration in latter stages. Sanctions for failing to meet these conditions include increased management, regression to previous stages, or ultimately revocation (with a return to mainstream prison). In Stage 1, participants are housed full-time in a secure environment. Programs include therapeutic intervention addressing dynamic risk factors for drug related offending,

adult education and work readiness programs, and prosocial living skills content. Abstinence is emphasised with gradual and medically assisted withdrawal, no contact visits, and frequent drug tests. A therapeutic prison environment is upheld through respectful staff relationships that support continuing abstinence, engage participants in treatment, and focus on prevention and pro-social modelling rather than rule violation detection. The likelihood of relapse is recognised and managed therapeutically rather than punished. In Stage 2, participants are able to leave the CDTCC to access community-based interventions. These interventions include employment, adult education and vocational training, volunteer work and social opportunities to assist in effective re-integration. In Stage 3, participants live outside the CDTCC facility but under intensive supervision. Programs in the community maintain and expand gains made in previous stages. Ongoing judicial supervision is a key feature of the program as the Judge oversees all decisions related to participant regression, removal, or progression, and regularly reviews Stage 2 and Stage 3 participants in person at the Drug Court. After completing the program, offenders are released into the community once their sentence period concludes. Therefore, to guard against any antitherapeutic effect of compulsory participation, treatment duration does not exceed the original sentence set by the sentencing court and participants have the added incentive of living in the community before their non-parole period expires.

The program follows other international examples of compulsory drug treatment, such as the United States' abstinence approach to drug use and the Dutch SOV regulation, which can impose up to two years of drug treatment detention for offenders considered a serious "nuisance element" (Oei, 2005). An evaluation study of the three-stage SOV compulsory program found that, soon after starting the program, respondents reported a substantial improvement of perceived physical and mental health and self-esteem which sustained after the

program ended. They also performed significantly better than offenders in the regular detention group in terms of subsequent offending, addiction and social functioning (Koeter & Bakker, 2007). Another stream of research on mostly voluntary Therapeutic Communities (TC) also shows some promising results in reducing drug use and recidivism outcomes (Mitchell et al., 2007; Mitchell et al., 2012; de Andrade et al., 2018; Doyle et al., 2019). TCs share similar characteristics with the CDTCC in that participants are housed separately full-time with treatment stages focusing on re-socialisation, intensive therapy, and gradually increasing responsibilities.

To our knowledge there has only been one evaluation of the NSW CDTCC. A study by Dekker and colleagues (2010) from the NSW Bureau of Crime Statistics and Research (BOCSAR) found improvements in mental and physical health and treatment readiness at Stages 1 and 2 compared to baseline. Further, only a very small proportion (1.8%) of drug tests were classified as 'positive,' although illegal and non-prescribed drugs were detected in at least one test for most participants and a greater proportion of drug tests were positive in Stage 3 compared to Stages 1 and 2. Contrary to the compulsory nature of the program, the vast majority of participants perceived their attendance as voluntary and felt positive about the program. While these findings are promising, one limitation of the study was the diminishing number of respondents across the stages, with 95 respondents before Stage 1 to only 13 by the end of Stage 3. The small sample sizes made it difficult to be sure the experiences of those assessed were representative of participants in general, and changes in interview data from baseline through to the end of Stage 3 could not be investigated.

Aims

The primary purpose of the current study was to evaluate whether participation in the CDTP was associated with change in risk factors that are relevant to the drug-crime cycle. Participants completed a battery of self-report psychometric measures before they started the program and at the end of each of the program stages to assess a range of key outcomes targeted by treatment including self-efficacy, quality of life, attitudes towards crime and offending, thinking styles that support criminality, problem solving ability, psychological and social functioning, and self-control. Key responsivity factors, including treatment motivation and readiness, and perceived coercion, were also measured before they started the program and at the end of each program stage. Responsivity factors may not be related to recidivism directly, but they can affect an individual's response to treatment and, thus, the efficacy of treatment (Andrews & Bonta, 2010). Therefore, information on these factors can assist with tailoring treatment to maximize participants' ability to learn from an intervention. We examined whether participants showed change on treatment targets and responsivity measures before they started the program and between treatment stages.

METHODS

Participants

Five hundred and thirty-eight male participants had entered the CDTCC at the time of analysis. To be included in our analyses, participants needed to have completed at least one assessment either before starting the program or after one of the program stages. One person was removed because they only completed one subscale on one measure at a single timepoint. Given that some participants completed some, but not all assessments, the final

sample size for each measure varied (see Table 1 for all sample sizes).

Measures

A battery of self-report psychometric measures was administered to participants at four timepoints, once before they started the program and at the end of each of the program stages. For the purposes of this study, we examined selected measures in the battery relevant specifically to targets of intervention and key participant responsivity factors.

Treatment targets

The program targets various specific, measurable aspects of participant functioning to improve over the course of treatment. Data was collected at each timepoint on some of these targets, namely, self-efficacy, quality of life, criminal attitudes, criminal thinking styles, problem solving, psychological and social functioning, and self-control.

Drug-taking Confidence Questionnaire (DTCQ). On the 50-item DTCQ (0 = not confident at all to 100 = very confident; Annis & Martin, 1985), participants rated how confident they were that they could resist using their nominated drug in high risk situations which have been demonstrated to be linked to drug use (when experiencing unpleasant emotions, physical discomfort, pleasant emotions, urges or temptations to use, conflict with others, social pressure to use, and pleasant times with others, and if they wanted to test personal control). Higher average confidence ratings indicate higher self-efficacy.

Quality of Life Inventory (QOLI). The 32-item QOLI (below 0 to -6 = increasing dissatisfaction; above 0 to 6 = increasing satisfaction; Frisch et al., 1992) measures satisfaction with sixteen physical, social and psychological aspects of life (health, self-esteem, goals and values, money, work, play, learning, creativity, helping, love, friends, children,

relatives, home, neighbourhood, and community). Higher scores indicate more satisfaction in the area.

Crime PICS II (CP-II). This tool consists of 20 items (1 = strongly agree to 5 = strongly disagree) and a 15-item problem inventory (1 = a big problem to 4 = no problem) to measure attitudes towards offending and crime (Frude et al., 2009). The measure provides a main score which represents a person's general attitudes to offending, as well as specific measures on: anticipation of re-offending, victim hurt denial, evaluation of crime as worthwhile, and perception of current life problems. High scores indicate the individual has attitudes which predispose him towards involvement in crime, or, in the case of the problems inventory, has problems in many areas of his life.

Psychological Inventory of Criminal Thinking Styles (PICTS). The 80-item PICTS (1 = disagree to 4 = strongly agree; Walters, 1995) assesses criminal thinking styles believed to support a criminal lifestyle (mollification, cutoff, entitlement, power orientation, sentimentality, superoptimism, cognitive indolence and discontinuity). It also includes two subscales used to assess response styles, which were not analysed as they may not be considered key targets of intervention. Higher ratings indicate participants exhibit thinking styles that may support criminality.

The Social Problem Solving Inventory-Revised: Short (SPSI-R:S). The 25-item SPSI-R:S (0 = not at all true to 4 = extremely true; D'Zurilla et al., 2002) assesses social problem solving ability. Raw scores are converted to standard scores so that total score and subscales score have an average of 100. The tool consists of five subscales. Higher ratings on the positive problem orientation and rational problem-solving style subscales, and lower ratings of negative problem orientation, impulsivity/carelessness style, and avoidance style subscales indicate good social problem solving.

The Self-Rating Form (SRF) - Psychological and Social Functioning domains. This 94-item tool (0 = strongly disagree to 4 = strongly agree; Simpson & Knight, 1998) measures psychosocial functioning and motivational factors in three domains which are divided into 13 subdomains. The two domains that were considered targets for intervention included psychological functioning (self-esteem, depression, anxiety, decision making, self-efficacy) and social functioning¹ (hostility, risk taking, social conformity). Higher scores indicate exhibiting more of that factor in each subdomain.

The Self Control Scale (SCS). The 24-item SCS (1 = strongly disagree to 4 = strongly agree; Grasmick et al., 1993) measures an individual's self-control in six domains: impulsivity, simple tasks, risk-seeking, physical activities, self-centeredness, and temper. Higher scores indicate more self-control.

Responsivity factors

Data was collected at each timepoint on participants' treatment motivation and readiness as well as perceived coercion to participate in the program.

The Self-Rating Form (SRF) - Treatment Motivation domain. The SRF (described above) measures treatment motivation which includes problem recognition, desire for help, treatment readiness, external pressures subdomains. Higher scores indicate exhibiting more of that factor in the subdomains.

Corrections Victoria Treatment Readiness Questionnaire (CVTRQ). The 20-item CVTRQ (1 = strongly disagree to 5 = strongly agree; Casey et al., 2007) assesses an offender's readiness for treatment programs. Higher scores indicate a higher degree of readiness to participate and engage in

¹ We excluded the 'childhood problems' subscale from our analyses as these scores "represent deviant attitudes and actions early in development" which were not targets of treatment (Knight et al., 2003, p. 55).

treatment. Offenders with a total score of 72 and above are classified as 'program ready'.

MacArthur Perceived Coercion Scale (MPCS). The 5-item MPCS (0 = yes, 1 = don't know, 2 = no; Gardner et al., 1993) assesses perceived coercion in participating in the program. Higher scores indicate greater perceived coercion to enter treatment.

General recidivism risk and needs

Level of Service Inventory-Revised (LSI-R). The 54-item LSI-R (Andrews & Bonta, 1995) is designed to classify an offender's risk of reoffending and identify their criminogenic needs. It consists of items that measure one static (unchangeable through intervention) and nine dynamic risk factors that are considered universally related to crime: Criminal History (static), Education/Employment, Finance, Family/Marital, Accommodation, Leisure/Recreation, Companions, Alcohol/Drug problems, Emotional/Personal problems, and Attitudes/Orientation. Higher scores indicate increased needs in that domain.

For the purposes of this study, LSI-R scores were examined only for the purposes of generating a profile of the characteristics of CDTP participants at baseline, and we did not assess change on this measure over the course of treatment.

Data analyses

Data on relevant variables were missing for a number of participants. In particular, not all participants completed the full battery of psychometric measures at the end of each stage. Further, obvious individual errors made in data entry were excluded from analysis and if more than two errors were made on a measure, responses for that measure were removed for that timepoint.

We ran mixed effects models to analyse the data for a number of reasons. A primary consideration was their ability to include all available data; as such, we were able to include all participants who completed a

measure at any timepoint rather than restricting the analysis to only those who completed the measure at all timepoints which would lead to biased and inefficient estimates. These models also allow for the inclusion of both fixed effects, which are model components used to define systematic relationships such as overall changes over time, and random effects, which account for variability among subjects (see e.g., Garcia & Marder, 2017 for a full explanation of possible statistical approaches for longitudinal data). For the purposes of assessing within-treatment change we ran mixed effects models analyses with program stage as the fixed factor, and participants as a random factor.² We looked at F tests results to determine whether responses on scales significantly differed between stages. If so, we examined post-hoc results to determine significant changes between stages.^{3, 4}

² We examined the interclass correlation coefficient (ICC) of each model to help determine whether participant should be set as a random factor. In this case, an ICC of zero or very close to zero means the observations within participant clusters are no more similar than observations from different clusters. ICCs in this study ranged from .13-.72, suggesting that mixed models were necessary.

³ Because a large number of analyses were run, thus potentially increasing the possibility of false positives, we predetermined that a p -value less than or equal to .01 (rather than .05) indicates a statistically significant result.

⁴ Analyses were rerun to ensure various factors did not impact results. First, we split participants into those who received the RUSH program (which was implemented from group 30 onwards) versus those who received precursor programs. Second, we split participants into those who had access to Methadone or Buprenorphine treatment if needed versus those who did not. Third, we excluded responses from participants who had completed the program a second time, retaining their initial responses. Fourth, we excluded participants who experienced COVID-19-related leave restrictions. Last, if participants regressed to a previous stage, thus completing the battery twice for that stage, we replaced their initial responses with responses given at regression (e.g., if they completed Stage 1 twice, we included their second set of responses). For all analyses, findings were similar to those found across all participants reported here, with few minor differences.

RESULTS

Needs and characteristics of CDTP participants at entry

Of those who completed the LSI-R prior to starting the program, the average assessed risk of recidivism was in the Medium category ($M = 33.75$, $SD = 6.56$). Only a small proportion were in the Low ($n = 5$; valid percent = 1.3%), Low-Medium ($n = 22$; valid percent = 5.6%), or High categories ($n = 48$; valid percent = 12.2%). The majority of participants were categorised as Medium ($n = 135$; 34.4%) or Medium-High ($n = 182$; 33.9%) risk.

Corrective Services NSW applies scoring thresholds to each of the domains of dynamic risk to categorise the severity of needs. These thresholds categorise scores to indicate 'strength', 'no immediate need for improvement', 'some need for improvement', and 'considerable need for improvement.' Scoring in the latter category suggests that the domain caused serious adjustment problems and contributed markedly to offending. On average, participants reported close to no needs on the Accommodation domain ($M = 1.29$, $SD = 1.05$), and some needs on the Education/Employment ($M = 7.28$, $SD = 2.07$), Family/Marital ($M = 1.80$, $SD = 1.22$), Companions ($M = 2.38$, $SD = .84$), Emotional/Personal ($M = 2.01$, $SD = 1.50$), and Attitudes/Orientation ($M = 2.04$, $SD = 1.36$) domains. They reported close to considerable needs on the Financial ($M = 1.73$, $SD = .52$) and Leisure/Recreation domains ($M = 1.74$, $SD = .59$). Of particular interest in our sample, participants reported considerable need for improvement in the Alcohol/Drug problems domain ($M = 6.14$, $SD = 1.46$). In total, 90.9% of those who completed the Alcohol/Drug domain assessment of the LSI-R reported considerable need for improvement on the domain.

In terms of participants' ratings of treatment targets on the psychometric battery prior to starting the program, they tended to rate themselves as average or better on many of the measures, including subscales of self-efficacy, problem solving ability, criminal attitudes, criminal thinking styles, and self-control measures (see Appendix 1). However, almost half of the participants reported having a very low quality of life, suggesting that they felt unhappy and dissatisfied with life.

In reference to responsivity factors at baseline, participants' treatment readiness (CVTRQ) was high on average before they started the program, with most (84.5%) already crossing the threshold for being treatment ready. Similarly, they rated highly on average on the treatment motivation subscales of the SRF. Perceived coercion on the MPCS and external pressures subscale of the SRF was also low on average before participants began the program. These results suggest that participants tended to be treatment ready and considered attendance voluntary at the time of commencing the CDTP.

Within-treatment change

There were significant effects of time across all measures on both total and subscale scores, except on the SCS physical activities subscale (see Appendix 1 for F and p values). In other words, responses on measures changed significantly across the program stages. Specific changes between each stage are broken down below, with the exception the SCS physical activities subscale which was excluded from follow-up analyses.

From baseline to Stage 1

Treatment Targets. First, we examined whether there were changes on treatment target outcomes before participants started the program compared to the end of Stage 1 of the program. Participants' confidence in their ability to resist drugs (DTCQ), quality of life (QOLI), psychological and social functioning (SRF), criminal attitudes (CP-II), thinking

styles (PICTS), problem solving ability (SPSI-R:S), and self-control (SCS) were significantly improved at every timepoint after they started the program compared to before they started the program (see Figures 1 to 8; see Appendix 1). In other words, they reported improvements on all measures after starting the program compared to before.

Responsivity factors. Participants also reported significant change on most responsivity measures. Treatment readiness on the CVTRQ improved at the end of Stage 1 compared to baseline (Figure 9). Scores on three of the subdomains within the treatment motivation domain of the SRF were lower at the end of Stage 1 compared to baseline, while the fourth exhibited no change, suggesting that problem recognition, desire for help, and external pressure decreased, while treatment readiness remained stable (Figure 11). Perceived coercion on the MPCS was lower at the end of Stage 1 compared to baseline (Figure 10).




Figure 1. Average total self-efficacy score (DTCQ)




Figure 2. Average overall quality of life score (QOLI)




Figure 3. Average psychological functioning scores




Figure 4. Average social functioning scores (SRF)




Figure 5. Average general attitudes to offending scores (CP-II)




Figure 6. Average total problem-solving score (SPSI-R:S)




Figure 7. Average thinking styles scores (PICTS)




Figure 8. Average self-control scores (SCS)

From Stage 1 to Stage 2

Treatment targets. On average, at the end of Stage 2, compared to at the end of Stage 1, participants reported significant change on aspects of all treatment target measures. They reported improved: 1) confidence in their ability to resist drugs across all high-risk situations and many of the subscales (when experiencing unpleasant emotions, urges or temptations to use, social pressure to use, pleasant times with others, and if they wanted to test personal control; DTCQ), 2) on overall quality of life and the work, children, relatives, home, neighbourhood, and community subscales (QOLI), 3) psychological functioning on the self-esteem, depression, and decision making subscales and social functioning on all subscales (SRF), 4) general attitudes to offending, evaluation of crime as worthwhile, and perception of current life problems (CP-II), 5) cut off, cognitive indolence, and discontinuous criminal thinking styles (PICTS), 6) impulsive/careless and avoidant problem-solving (SPSI-R:S), and 7) impulsiveness and risk-taking (SCS).

Responsivity factors. Participants reported decreased treatment motivation on all subdomains of the SRF measure at the end of Stage 2 compared to at the end of Stage 1. There was no significant change on treatment readiness (TRQ) and perceived coercion (MPCS).

From Stage 2 to Stage 3

Treatment targets. On average, at the end of the program, compared to at the end of the previous stage, participants reported significant change on aspects of all treatment measures except self-efficacy and problem-solving. Specifically, they reported improvements in: 1) overall quality of life and the money, work, and home subscales (QOLI), 2) psychological functioning on the self-esteem, depression, and anxiety subscales as well as social functioning on the risk-taking subscale (SRF), 3) general attitudes to offending and perception of

current life problems (CP-II), 4) cognitive indolence and discontinuous thinking styles (PICTS), and 5) and impulsiveness (SCS).

Responsivity factors. Participants reported decreased treatment motivation on all subdomains (SRF) at the end of Stage 3 compared to Stage 2. There was no significant change in treatment readiness (TRQ) and perceived coercion (MPCS).




Figure 9. Average treatment readiness (CVTRQ) scores




Figure 10. Average perceived coercion scores (MPCS)




Figure 11. Average treatment motivation (SRF) scores.

DISCUSSION

With the widespread prevalence of drug use among people who are involved in the criminal justice system and its significant impact on the risk of reoffending, it is imperative to evaluate the CDT model as a major judicial avenue for providing intervention to address substance use-related needs. Consistent with RNR principles (e.g., Andrews & Bonta, 2010), which highlights the importance of treatment influencing dynamic risk factors to then reduce reoffending, our primary aim of this study was to investigate whether CDT participants exhibit change in a range of risk factors over the course of treatment. We also aimed to examine the nature of, and change in, responsivity factors that may affect the efficacy of treatment.

Participant needs and characteristics

Interestingly, participants tended to rate themselves as average or above average on a number of treatment target measures before they had begun treatment. One potential explanation is that some measures chosen to examine improvements from treatment may reflect dynamic risk factors that are highly heterogeneous across CDT participants or not commonly experienced by a number of participants. Alternatively, the results may indicate that some participants were unable or unwilling to accurately report on their own needs. While tendencies towards socially desirable responding are a recognised challenge in assessing needs among people involved in the criminal justice system (e.g., Juarez & Howard, 2018), there may also be clinical implications. For example, we found that on average, participants expressed at least a modest level of confidence in their ability to resist drugs at baseline. This may be of concern because initial preliminary data from the centre found that participants who were confident were more likely to relapse. Indeed, research suggests that high self-reported self-efficacy may indicate denial or

overconfidence which could, in turn, result in negative consequences such as making less effort to acquire the skills necessary to cope with problem behaviour (Burling et al., 1989).

Notwithstanding the above observations, our results indicated that CDTP participants tended to be of relatively high risk of general recidivism with a complex range of criminogenic needs (as assessed by the LSI-R), demonstrating the potential value of prioritising intensive intervention for this cohort. Despite their needs, participants gave good ratings of treatment readiness and other indicators of responsivity at the time of commencing the program, on average. This is consistent with the prior evaluation (Dekker et al., 2010) and provides promising indications that participants tended to be treatment ready and perceived a low level of coercion associated with their engagement in the program.

Change in treatment targets

Overall, we expected participants to improve on treatment target outcomes over the course of treatment. Indeed, our results gave promising initial indications that redirecting drug-dependent offenders into compulsory treatment and rehabilitation was successful in producing measurable change in a range of outcomes once they started the program. Specifically, global effects of participation in the CDTP included significantly improved self-efficacy, psychological and social functioning, criminal attitudes, criminal thinking styles, problem solving ability, and self-control from before starting the program to after. These improvements could be promising for more direct drug-related outcomes such as abstinence (e.g., Chavarria et al., 2012; Ilgen et al., 2005).

Our analyses also gave some insights into the incremental effects of each of the CDTP stages on changes in assessed treatment targets. We found that aspects of all measures further improved during Stage 2, and again during Stage 3 with the

exception of self-efficacy and problem-solving ability. Self-efficacy peaked at the end of Stage 1 before plateauing, possibly because participants' scores were already very high by that timepoint, leaving little room to improve further. Problem-solving ability, on the other hand, remained around average at all timepoints and did not improve after Stage 2, suggesting that treatment did not influence this construct to a great extent, especially at the final stage. Overall, however, there was value to every stage of the program with outcomes changing in a positive direction throughout.

Change in responsivity factors

Studies have shown that people who receive treatment ordered or supervised by the criminal justice system perceive greater external pressure to be in treatment. The literature is mixed on the impacts of compulsory treatment on substance use compared with control interventions (e.g., Werb et al., 2016). Despite the involuntary nature of the CDTP, however, coercion scores were low even before starting the program and decreased further by the end. Treatment readiness was also high before participants started the program, and increased further after they started. Therefore, participants appeared to consider attendance as voluntary and wanted to reduce their drug use. This desire to engage with treatment is associated with indicators of therapeutic engagement (Hiller et al., 2002) which, in turn, is related to greater likelihood of change after treatment (Garnick et al., 2012). As such, tailoring treatment to continue maintaining or improving these responsivity factors is highly likely to be beneficial to the success of the CDTP model.

We note seemingly contradictory findings in treatment readiness on the CVTRQ and treatment motivation on the SRF. This may be accounted for by the nature of the assessment items and underlying constructs. The CVTRQ mostly focuses on how respondents feel about treatment in general (e.g., "treatment programmes don't work") and their

own past offending (e.g., “I feel guilty about my offending”) which may be expected to improve or remain high over the course of treatment. Treatment motivation on the SRF, on the other hand, measures current feelings on problem recognition, desire for help, treatment readiness, and external pressures which may naturally diminish over time as participants experience therapeutic improvements (e.g., “In your opinion, your drug use is a problem for you”; “You need help in dealing with your drug use”). This scale of the SRF also includes items that appear to be context-specific or difficult for participants to interpret or rate under certain circumstances (e.g., “It is urgent that you find help immediately for your drug use” among those who have entered the program; “You plan to stay in this treatment program for a while” among those who are completing Stage 3), which has implications for its utility in assessing within-treatment change in motivation for CDTP participants.

Limitations

This study has limitations that should be considered. Significantly, we did not obtain psychometric ratings for an equivalent comparison group who did not participate in the program. Without a comparison group, we cannot conclude whether the observed changes can be attributed to the program or simply reflect spontaneous change over time. Given the self-report nature of the measures, there is also a possibility that participants may have provided responses that presented themselves in a favourable light. Although the extent of socially desirable responding in offenders is smaller than assumed (Mathie & Wakeling, 2011; Juarez & Howard, 2022), research has shown that inmates’ self-report responses can be impacted by changing context, with underreporting of risk factors becoming more likely at post-treatment compared to pre-treatment (Juarez & Howard, 2018). Therefore, reported

reductions in negative thoughts and behaviours may not reflect true change.

A related limitation is that our analyses did not include assessments of reoffending outcomes. In the context of evaluating within-treatment change, associations between measures of interest and recidivism give important information about the predictive validity of psychometric assessments and the extent to which underlying constructs represent key treatment targets. Without this analysis, the mechanisms of change in treatment remain unclear, along with whether within-treatment change reflects or influences significant post-release behaviours. This is especially so given that some measures in this study have limited empirical support for their predictive validity and change in some underlying constructs have not been found to be consistently related to reoffending (e.g., Banse et al., 2013; Hiller et al., 2006; Juarez & Howard, 2018). We intend to conduct additional research in the future to examine relationships between patterns of within-treatment change observed in the current study and recidivism outcomes, in order to generate additional insights about the therapeutic gains and outcomes associated with participating in the CDTP.

Conclusions

Overall, this study provides initial indications that participation in the CDTP may be a promising response to persistent drug-related crime, given the improvements we found in measured treatment targets that are expected to have an influence on drug-related offending. Improvements were often incremental and significant across each of the three program stages, suggesting additional benefits of every component of the CDTP. It is possible that the extent of change on treatment targets observed among participants was facilitated by their responsivity to the program, which was assessed as positive at baseline and continued to improve over the course of treatment; unfortunately, it was

beyond the scope of this study to conduct comprehensive analyses on interactions between responsiveness and magnitude of within-treatment change. Nonetheless, the findings of this study are one of the first to contribute to an understanding of intermediate processes of change associated with participation in the CDTCC, and will be supported by additional evaluations in the future that further investigate program processes and outcomes.

REFERENCES

- Andrews, D. A., & Bonta, J. (1995). *The Level of Service Inventory – Revised*. Toronto: Multi-Health Systems.
- Andrews, D. A., & Bonta, J. (2010). Rehabilitating criminal justice policy and practice. *Psychology, Public Policy, and Law, 16*(1), 39–55.
<https://doi.org/10.1037/a0018362>
- Annis, H.M., Sklar, S.M. & Turner, N.E. (1997). *The Drug-Taking Confidence Questionnaire (DTCQ): User's Guide*. Toronto, Canada: Addiction Research Foundation, Centre for Addiction and Mental Health.
- Australian Institute of Health and Welfare (2019). *The health of Australia's prisoners 2018*. Cat. no. PHE 246. Canberra: AIHW.
- Banse R., Koppehele-Gossel J., Kistemaker L. M., Werner L. A., Schmidt A. F. (2013). Pro-criminal attitudes, intervention, and recidivism. *Aggression and Violent Behavior, 18*(6), 673–685.
<https://doi.org/10.1016/j.avb.2013.07.024>
- Bennett, Trevor & Holloway, Katy & Farrington, David. (2008). The Statistical Association between Drug Misuse and Crime: A Meta-analysis. *Aggression and Violent Behavior, 13*(2), 107–118.
<https://doi.org/10.1016/j.avb.2008.02.001>
- Birgden, A., & Grant, L. (2010). Establishing a compulsory drug treatment prison: Therapeutic policy, principles, and practices in addressing offender rights and rehabilitation. *International journal of law and psychiatry, 33*(5–6), 341–349.
<https://doi.org/10.1016/j.ijlp.2010.09.006>
- Burling, T. A., Reilly, P. M., Moltzen, J. O., & Ziff, D. C. (1989). Self-efficacy and relapse among inpatient drug and alcohol abusers: a predictor of outcome. *Journal of Studies on Alcohol, 50*, 354–360.
<https://doi.org/10.15288/jsa.1989.50.354>
- Casey, S., Day, A., Howells, K., & Ward., T. (2007). Assessing suitability for offender rehabilitation: Development and validation of the Treatment Readiness Questionnaire. *Criminal Justice and Behaviour, 34*(11), 1427–1440.
<https://doi.org/10.1177/0093854807305827>
- Chavarria, J., Stevens, E. B., Jason, L. A., & Ferrari, J. R. (2012). The Effects of Self-Regulation and Self-Efficacy on Substance Use Abstinence. *Alcoholism treatment quarterly, 30*(4), 422–432.
<https://doi.org/10.1080/07347324.2012.718960>
- Crimes (Administration of Sentences) Act 1999* (NSW), s 106B.
<https://legislation.nsw.gov.au/view/html/inforce/current/act-1999-093#sec.106B>
- de Andrade, D., Ritchie, J., Rowlands, M., Mann, E., & Hides, L. (2018). Substance Use and Recidivism Outcomes for Prison-Based Drug and Alcohol Interventions. *Epidemiologic reviews, 40*(1), 121–133.
<https://doi.org/10.1093/epirev/mxy004>
- Dekker, J., Smith, N., & O'Brien, K. (2010). *An evaluation of the compulsory drug treatment program (CDTP)*. Department of Justice and Attorney General, NSW Bureau of Crime Statistics and Research. <https://www.bocsar.nsw.gov.au/Publications/Legislative/120.pdf>
- Dowden, C., & Brown, S. L. (2002). The role of substance abuse factors in predicting recidivism: A meta-analysis. *Psychology, Crime & Law, 8*(3), 243–264. <https://doi.org/10.1080/10683160208401818>
- Doyle, M.F., Shakeshaft, A., Guthrie, J., Snijder, M. and Butler, T. (2019), A systematic review of evaluations of prison-based alcohol and other drug use behavioural treatment for men. *Australian and New Zealand Journal of Public Health, 43*(2), 120–130.
<https://doi.org/10.1111/1753-6405.12884>
- D’Zurilla, T. J., Nezu, A. M., & Maydue-Olivares, A. (2002). *Social Problem Solving Inventory– Revised*. MHS
- Frisch, M., Cornell, J., Villanueva, M., & Retzlaff, P. (1992). Clinical validation of the Quality of Life Inventory. A measure of life satisfaction for use in treatment planning and outcome assessment. *Psychological Assessment, 4*(1), 92–101.
<https://doi.org/10.1037/1040-3590.4.1.92>
- Frude, N., Honess, T., & Maguire, M. (2009). *CRIME PICS–II Manual*. London: M&A Research. www.crime-pics.co.uk/cpicsmanual.pdf
- Garcia, T. P., & Marder, K. (2017). Statistical Approaches to Longitudinal Data Analysis in Neurodegenerative

- Diseases: Huntington's Disease as a Model. *Current neurology and neuroscience reports*, 17(2), 14. <https://doi.org/10.1007/s11910-017-0723-4>
- Gardner, W., Hoge, S., Bennett, N., Roth, L., Lidz, C., Monahan, J., & Mulvey, E. (1993). Two scales for measuring patients' perceptions of coercion during mental hospital admission. *Behavioural Sciences and the Law*, 11, 307-322. <https://doi.org/10.1002/bsl.2370110308>
- Garnick, D. W., Lee, M. T., O'Brien, P. L., Panas, L., Ritter, G. A., Acevedo, A., Garner, B. R., Funk, R. R., & Godley, M. D. (2012). The Washington circle engagement performance measures' association with adolescent treatment outcomes. *Drug and alcohol dependence*, 124(3), 250-258. <https://doi.org/10.1016/j.drugalcdep.2012.01.011>
- Grasmick, H. G., Tittle, C. R., Bursik, Jr, R. J., & Arneklev, B. J. (1993). Testing the core empirical implications of Gottfredson & Hirschi's general theory of crime. *Journal of Research in Crime and Delinquency*, 30, 5-29. <https://doi.org/10.1177/00224278930300010>
- Hiller, M. L., Knight, K., Leukefeld, C., & Simpson, D. D. (2002). Motivation as a predictor of therapeutic engagement in mandated residential substance abuse treatment. *Criminal Justice and Behavior*, 29(1), 56-75. <https://doi.org/10.1177/0093854802029001004>
- Hiller, M. L., Knight, K., Saum, C. A., & Simpson, D. D. (2006). Social Functioning, Treatment Dropout, and Recidivism of Probationers Mandated to a Modified Therapeutic Community. *Criminal Justice and Behavior*, 33(6), 738-759. <https://doi.org/10.1177/0093854806288242>
- Ilgen, M., McKellar, J., & Tiet, Q. (2005). Abstinence self-efficacy and abstinence 1 year after substance use disorder treatment. *Journal of Consulting and Clinical Psychology*, 73, 1157-1180. <https://doi.org/10.1037/0022-006X.73.6.1175>
- Juarez, T., & Howard, M. V. A. (2018). *Assessing offender change over treatment: The influence of treatment context on self-reported antisocial attitudes*. Sydney: Corrections Research Evaluation and Statistics, Corrective Services NSW.
- Juarez, T., & Howard, M. V. A. (2022). Self-reported change in antisocial attitudes and reoffending among a sample of 2,337 males convicted of violent offenses. *Criminal Justice and Behavior*, 49, 3-19. <https://doi.org/10.1177/00938548211013576>
- Knight, K., Simpson, D. D., & Hiller, M. L. (2003). *Outcome Assessment of Correctional Treatment (OACT)*. Texas Christian University Institute of Behavioral Research.
- Koeter, M. W. J., & Bakker, M. (2007). *E Effectevaluatie van de Strafrechtelijke Opvang Verslaafden (SOV)*. Retrieved from the Wetenschappelijk Onderzoek- en Documentatiecentrum, Ministry of Justice and Security: <https://repository.wodc.nl/handle/20.500.12832/1207>
- Mathie, N. L., & Wakeling, H. C. (2011). Assessing socially desirable responding and its impact on self-report measures among sexual offenders. *Psychology, Crime & Law*, 17, 215-237. <https://doi.org/10.1080/10683160903113681>
- Mitchell, O., Wilson, D. B., & MacKenzie, D. L. (2007). Does incarceration-based drug treatment reduce recidivism? A meta-analytic synthesis of the research. *Journal of Experimental Criminology*, 3(4), 353-375. <https://doi.org/10.1007/s11292-007-9040-2>
- Mitchell, O., Wilson, D.B. and MacKenzie, D.L. (2012). The Effectiveness of Incarceration-Based Drug Treatment on Criminal Behavior: A Systematic Review. *Campbell Systematic Reviews*, 8, i-76. <https://doi.org/10.4073/csr.2012.18>
- Oei, T. I. (2005). The SOV Regulation. A new criminal code regulation measured against the behavioral sciences. *European Journal of Psychiatry*, 19, 69-77. <https://doi.org/10.4321/S0213-61632005000200001>
- Simpson, D. D., & Knight, K. (1998). *The Texas Christian University Self Rating Form*. Texas Christian University. <http://ibr.tcu.edu/wp-content/uploads/2013/10/kk6-srf-95.pdf>
- Voce, A., & Sullivan, T. (2020). Drug use monitoring in Australia: Drug use among police detainees, 2020 (Statistical report no. 35). Australian Institute of Criminology.
- Walters, G. D. (2006). *The Psychology Inventory of Criminal Thinking Styles (PICTS) professional manual*. Allentown, PA: Center for Lifestyle Studies.
- Ward, T. (2002). Good lives and the rehabilitation of offenders: Promises and problems. *Aggression and Violent Behavior*, 7(5), 513-528.
- Werb, D., Kamarulzaman, A., Meacham, M. C., Rafful, C., Fischer, B., Strathdee, S. A., & Wood, E. (2016). The effectiveness of compulsory drug treatment: A systematic review. *The International journal on drug policy*, 28, 1-9. <https://doi.org/10.1016/j.drugpo.2015.12.005>

APPENDIX A

Estimated marginal means for psychometric measures at each point of assessment

Measure	Global model			Pre-S1		Post-S1		Post-S2		Post-S3		Significance (<i>p</i> _{Bonferroni})		
	<i>F</i>	<i>p</i>	<i>n</i>	M (SE)	<i>n</i>	M (SE)	<i>n</i>	M (SE)	<i>n</i>	M (SE)	Pre to S1	S1 to S2	S2 to S3	
Drug Taking Confidence Questionnaire (DTCQ)														
Total score	373.0	**	535	58.7 (0.79)	474	86.0 (0.83)	302	91.5 (1.03)	133	91.2 (1.53)	**	**	-	
Unpleasant emotions	399.0	**	535	53.0 (0.87)	474	83.9 (0.92)	302	89.9 (1.13)	133	90.7 (1.68)	**	**	-	
Physical discomfort	342.0	**	535	60.4 (0.82)	474	88.8 (0.86)	302	92.7 (1.07)	133	92.6 (1.60)	**	-	-	
Positive emotions	161.0	**	535	72.9 (0.78)	474	91.5 (0.82)	302	94.8 (1.02)	133	94.2 (1.53)	**	-	-	
Testing personal control	253.0	**	535	55.9 (0.99)	474	82.2 (1.04)	302	90.1 (1.29)	133	88.2 (1.89)	**	**	-	
Urges/temptations to use	343.0	**	535	55.7 (0.88)	474	84.5 (0.93)	302	90.7 (1.15)	133	90.8 (1.70)	**	**	-	
Conflict with others	274.0	**	535	63.3 (0.82)	473	88.8 (0.87)	302	92.6 (1.08)	133	92.9 (1.61)	**	-	-	
Social pressure to use	330.0	**	535	53.8 (0.94)	473	84.2 (1.00)	302	90.2 (1.23)	133	89.9 (1.82)	**	**	-	
Pleasant times with others	339.0	**	535	54.8 (0.92)	471	84.9 (0.97)	300	90.6 (1.20)	133	90.3 (1.77)	**	**	-	
Quality of Life Inventory (QOLI)														
Overall	131.0	**	533	0.90 (.04)	467	1.46 (.04)	303	1.77 (.05)	135	2.15 (.08)	**	**	**	
Health	38.5	**	532	1.69 (.11)	466	2.81 (.12)	301	2.95 (.14)	133	3.11 (.20)	**	-	-	
Self-esteem	87.8	**	532	1.40 (.10)	466	2.92 (.10)	302	3.03 (.13)	133	3.58 (.18)	**	-	-	
Goals	149.0	**	532	0.97 (.10)	465	3.18 (.11)	302	3.39 (.13)	133	3.87 (.19)	**	-	-	
Money	58.4	**	531	-0.84 (.11)	465	0.07 (.11)	302	0.54 (.14)	133	1.59 (.20)	**	-	**	
Work	98.3	**	531	-1.51 (.14)	466	-0.73 (.15)	302	1.00 (.18)	133	2.87 (.27)	**	**	**	
Play	50.3	**	532	0.76 (.10)	466	1.90 (.11)	301	2.13 (.13)	133	2.66 (.19)	**	-	-	
Learning	57.2	**	532	0.98 (.10)	466	2.37 (.10)	301	2.36 (.12)	132	2.53 (.18)	**	-	-	
Creativity	48.6	**	531	0.81 (.09)	466	1.83 (.09)	302	1.86 (.11)	132	2.31 (.16)	**	-	-	
Helping	61.5	**	532	1.11 (.10)	466	2.20 (.10)	302	2.56 (.12)	133	3.11 (.18)	**	-	-	
Love	39.7	**	531	0.78 (.14)	466	1.88 (.15)	301	2.36 (.18)	133	3.24 (.26)	**	-	-	
Friends	30.2	**	532	0.99 (.11)	466	1.96 (.11)	300	1.98 (.14)	133	2.48 (.20)	**	-	-	
Children	35.1	**	529	0.56 (.15)	466	1.29 (.16)	299	2.12 (.19)	132	2.97 (.27)	**	**	-	
Relatives	34.9	**	532	1.80 (.12)	466	2.65 (.13)	301	3.31 (.15)	133	3.27 (.22)	**	**	-	
Home	62.1	**	531	0.48 (.15)	466	1.38 (.16)	301	2.47 (.19)	133	3.74 (.27)	**	**	**	
Neighbourhood	30.4	**	532	0.22 (.11)	465	0.82 (.11)	301	1.33 (.14)	133	1.90 (.20)	**	*	-	
Community	48.9	**	530	0.42 (.10)	464	1.23 (.10)	302	1.79 (.13)	133	2.41 (.19)	**	*	-	
The Self-Rating Form (SRF)														
Psychological functioning														
Self-esteem	257.0	**	533	40.5 (.48)	475	51.1 (.50)	301	54.7 (.60)	133	58.9 (.86)	**	**	**	
Depression	135.0	**	533	31.0 (.40)	475	24.8 (.42)	304	23.1 (.50)	133	20.0 (.71)	**	*	**	
Anxiety	136.0	**	533	37.4 (.50)	475	29.7 (.53)	304	27.9 (.62)	133	24.0 (.87)	**	-	**	
Decision making	223.0	**	533	44.2 (.39)	475	53.0 (.41)	304	55.2 (.49)	133	56.8 (.71)	**	**	-	
Self-efficacy	73.9	**	533	52.6 (.39)	475	57.9 (.41)	303	58.5 (.50)	133	60.5 (.72)	**	-	-	
Social functioning														
Hostility	80.7	**	532	32.3 (.47)	475	27.6 (.49)	304	25.7 (.56)	133	23.5 (.77)	**	*	-	
Risk-taking	180.0	**	533	44.9 (.45)	475	38.1 (.48)	304	34.1 (.56)	133	30.5 (.79)	**	**	**	
Social conformity	97.9	**	533	48.0 (.31)	474	52.2 (.33)	303	53.5 (.38)	133	54.0 (.54)	**	*	-	
Treatment motivation														
Problem recognition	190.0	**	532	59.2 (.55)	474	53.0 (.58)	304	44.4 (.71)	133	36.7 (1.05)	**	**	**	
Desire to help	163.0	**	533	60.5 (.37)	475	56.9 (.39)	304	52.6 (.47)	133	46.7 (.68)	**	**	**	
Treatment readiness	33.3	**	533	60.7 (.35)	475	61.4 (.36)	304	59.4 (.44)	133	55.0 (.63)	-	**	**	
External pressure	93.6	**	533	46.7 (.45)	472	42.9 (.47)	304	39.9 (.57)	133	33.6 (.82)	**	**	**	
Crime PICS-II (CP-II)														
General attitude to offending	278.0	**	528	46.0 (0.41)	474	35.6 (0.43)	304	33.7 (0.51)	136	30.5 (0.73)	**	*	**	
Anticipation of re-offending	179.0	**	529	15.8 (0.16)	474	12.3 (0.17)	305	11.6 (0.21)	136	10.6 (0.30)	**	-	-	
Victim hurt denial	35.1	**	528	5.50 (0.09)	474	4.49 (0.10)	302	4.53 (0.12)	135	4.63 (0.17)	**	-	-	
Crime as worthwhile	170.1	**	529	11.8 (0.14)	474	9.12 (0.15)	305	8.48 (0.18)	136	7.86 (0.25)	**	*	-	
Current life problems	315.0	**	521	35.8 (.34)	468	29.1 (.35)	294	25.3 (.42)	132	21.0 (.59)	**	**	**	

Note. ** *p* < .001; * *p* < .01

(table continues)

Estimated marginal means for psychometric measures at each point of assessment

Psychological Inventory of Criminal Thinking Styles (PICTS)													
Mollification	77.4	**	530	14.8 (.17)	474	12.5 (.18)	300	12.4 (.21)	133	12.2 (.29)	**	-	-
Cutoff	221.0	**	529	19.3 (.18)	474	15.5 (.19)	302	14.3 (.23)	131	13.3 (.33)	**	**	-
Entitlement	50.7	**	528	14.5 (.15)	475	12.9 (.16)	301	12.7 (.19)	133	12.1 (.27)	**	-	-
Power orientation	65.5	**	529	14.0 (.16)	474	12.0 (.17)	301	11.6 (.20)	133	11.5 (.29)	**	-	-
Sentimentality	40.4	**	530	18.1 (.15)	474	17.0 (.16)	301	16.5 (.19)	133	15.9 (.26)	**	-	-
Superoptimism	19.8	**	529	16.9 (.18)	475	15.8 (.19)	302	15.6 (.22)	132	15.3 (.30)	**	-	-
Cognitive indolence	248.0	**	529	20.2 (.19)	475	15.8 (.20)	299	15.0 (.24)	133	13.8 (.34)	**	*	*
Discontinuity	227.0	**	530	20.4 (.18)	474	16.7 (.19)	302	15.8 (.22)	133	14.8 (.31)	**	**	*
The Social Problem Solving Inventory-Revised: Short (SPSI-R:S)													
Total	130.0	**	533	91.2 (.69)	463	103.1 (.74)	301	105.1 (.88)	131	108.9 (1.25)	**	-	-
Positive problem orientation	38.8	**	533	92.4 (.79)	466	100.7 (.85)	302	102.1 (1.04)	133	105.4 (1.53)	**	-	-
Negative problem orientation	41.5	**	533	99.2 (.67)	466	93.1 (.70)	302	91.6 (.83)	134	89.9 (1.17)	**	-	-
Rational problem solving	61.1	**	533	87.4 (.81)	466	97.5 (.86)	302	99.1 (1.05)	135	103.9 (1.53)	**	-	-
Impulsivity/Carelessness style	61.0	**	532	108.4 (.77)	466	100.1 (.82)	302	96.5 (.98)	133	95.2 (1.4)	**	*	-
Avoidance	51.4	**	533	105.0 (.68)	466	98.6 (.72)	302	95.3 (.86)	134	94.1 (1.23)	**	*	-
Self Control Scale (SCS)													
Impulsivity	207.0	**	531	9.29 (.09)	474	11.21 (.09)	305	11.78 (.11)	137	12.67 (.17)	**	**	**
Simple tasks	76.0	**	531	9.85 (.08)	475	10.84 (.09)	305	11.11 (.10)	138	11.49 (.14)	**	-	-
Risk-taking	79.3	**	531	10.00 (.08)	475	11.01 (.08)	305	11.62 (.10)	137	11.66 (.15)	**	**	-
Physical activities	2.03	-	531	8.86 (.08)	475	8.79 (.08)	305	8.72 (.09)	138	9.06 (.13)	NA	NA	NA
Self-centredness	24.8	**	531	11.4 (.08)	475	11.9 (.08)	304	12.1 (.10)	137	12.3 (.14)	**	-	-
Temper	45.3	**	531	11.2 (.10)	475	12.2 (.10)	305	12.5 (.12)	138	12.5 (.17)	**	-	-
Corrections Victoria	55.8	**	387	79.7 (0.41)	341	84.5 (0.43)	222	85.8 (0.52)	121	84.7 (0.67)	**	-	-
Treatment Readiness Questionnaire (CVTRQ)													
MacArthur Perceived Coercion Scale (MPCS)	32.6	**	477	2.37 (0.10)	335	1.45 (0.11)	221	1.33 (0.13)	127	1.09 (0.16)	**	-	-

Note. ** $p < .001$; * $p < .01$

Other CRES

Research Titles

Dec 2023 Assessment and experience of prison climate in NSW correctional centres

Dec 2023 Five Minute Intervention (FMI) skills acquisition by correctional staff: The role of manager buy-in

Dec 2023 Dynamics of wellbeing and needs satisfaction among people in prison

Dec 2023 Empirical review of the Pathways to Employment (P2E) pilot for women in prison

Dec 2023 Conducting survey research using inmate digital tablets: Lessons learned from research conducted in NSW correctional centres

Oct 2023 Assessing the impact of Five Minute Interventions (FMI) training on behavioural indicators of correctional centre safety and order: An interrupted time series analysis

Oct 2023 Impacts of digital tablets on trends in correctional centre safety and order: A controlled interrupted time series study

Sept 2023 Development and validation of the Triage Risk Assessment Scale: Sexual Offending (TRAS:SO)

Sept 2023 Review of the birth certificate Identity Management Enhancement pilot

July 2023 Implementation of an initiative for people with less than five months to serve in prison: Staff insights on the Short Sentence Intensive Program (SSIP)

June 2023 The Intensive Learning Centre: Staff experiences of implementation and the SILC reforms

April 2023 Five Minute Interventions (FMI): Custodial staff views and experiences of implementing FMI

Dec 2022 Effects of the Workplace Mentor Program on correctional officers' perceptions of workplace culture

Dec 2022 Five Minute Interventions (FMI): Long-term effects of training on custodial staff attitudes towards prisoners, motivation and ability to support rehabilitation, and job stress and satisfaction

Nov 2022 Effects of Words @ Work training on perceptions of offender rehabilitation and job experiences among Corrective Services Industries (CSI) overseers

Sept 2022 Offender participation outcomes and predictors of treatment completion in the High Intensity Program Units (HIPUs)

Aug 2022 Staff and inmate experiences of prison social climate at Rapid Build correctional centres: A quantitative evaluation

July 2022 Evaluation of the Alternate Sanctions Program (ASP): Within-treatment Change

June 2022 Implementing digital technologies in prisons: Inmate uptake and perceived value of in-cell digital tablets

April 2022 Actuarial assessment of domestic violence recidivism risk among custody-based males: The Domestic Violence – Triage Risk Assessment Scale (DV-TRAS)

March 2022 A qualitative exploration of factors influencing prison social climate at Rapid Build and traditional correctional centres

Jan 2022 Quality of the therapeutic alliance and associations with program outcomes for offenders participating in High Intensity Program Units (HIPUs)

Dec 2021 The Initial Transitional Support (ITS) program: Implementation evaluation

Dec 2021 How does the role of custodial staff influence their perceptions of offender rehabilitation and responses to Five Minute Interventions (FMI) training?

Oct 2021 Five Minute Interventions (FMI): Short-term effects of training on staff attitudes towards prisoners, motivation and ability to support rehabilitation, and job stress and satisfaction



Research Bulletin No.63
ISSN 2207 0850
© Corrective Services NSW

Corrections Research, Evaluation & Statistics
Delivery, Performance & Culture
Corrective Services NSW
GPO Box 31
Sydney NSW Australia
Telephone: (02) 8346 1556
Email: research.enquiries@dcj.nsw.gov.au